

Greengrazer V

SORGHUM-SUDANGRASS

(*Sorghum bicolor x Sorghum Sudanese*)

- Small-seeded and thin-stemmed type
- Dark green color
- Anthracnose and Downy Mildew resistant

Greengrazer V is a small-seeded three way cross with thin stems that are highly palatable. Regrowth after cutting is very fast. Planting Greengrazer V at higher populations per acre will result in a finer stemmed forage. Finer stems will allow the forage to dry faster for higher quality hay than is possible with thick stemmed types. For grazing, wait until plants are 18"-36" tall. Remove cattle when plant height has been grazed down to 8".

Disease Ratings:

Anthracnose: R
Downy Mildew: R

Agronomic Traits:

Early Seedling Vigor: Excellent
Growth Habit: Upright
Recovery After Cutting: Excellent
Maturity: 60 days to boot
Uniformity: Excellent
Plant Type: Juicy Stalk
Midrib Type: Conventional

Planting Rates:

Approx. Seeds Per Pound: 19,500
Rate (Lbs.) **Dryland** **Irrigated**
Broadcast: 35-50 50-65
Drill: 30-45 45-60

Adaptation Ratings:

Photosynthetic Type: C4 - Warm Season
Soil Temperature: Warm (60 F)
Water Requirement: Low

Seeding:

- Soil temperature should be a minimum of 62 F.
- Can be no-tilled into the stubble of winter and spring crops.
- Planting depth should be 1".
- Do not plant in soils with pH greater than 7.5.

Harvest:

- Greengrazer V is usually harvested 60-65 days after seeding.
- Protein will decline as harvest is delayed, but energy will increase upon heading due to continued sugar formation in the sorghum stalks and leaves, and carbohydrate deposition in the developing grains.

Crop Use Information:

Life Cycle: Annual
Ease of Establishment: Good
Shade Tolerance: Poor - Fair
Drought Stress: Excellent
Wet Soil: Fair
Low pH Tolerance: Moderate
Minimum pH: 6.0
Saline Soils (White Alkali): Fair
Saline - Sodic Soils (Black Alkali): Poor - Fair
Hay: Excellent
Silage: Excellent
Continuous Grazing: Do not Continuous Graze
Rotational Grazing: Excellent
Palatability: Excellent

Strengths

- High yield potential.
- Thin-stemmed plant type.
- Small-seeded type.
- Dark green plant color.
- Good overall disease resistance package.
- Resistant to Anthracnose and Downy Mildew.
- Sweet juicy midrib.

Avoid Nitrate and Prussic Acid Poisoning from sorghum:
Avoid large nitrogen applications prior to expected drought periods.

2,4-D can increase Prussic Acid concentration for several weeks after application. Do not harvest drought-damaged plants within 4 days following a good rain.

Do not green chop within 7 days of a killing frost. Cut at a higher stubble height, nitrates tend to accumulate in the lower stalk. Wait 1 month before feeding silage to give Prussic Acid enough time to dissipate.

